



Loss of labile organic carbon from subsoil due to land-use changes in subtropical China



Hao Sheng^{a,*}, Ping Zhou^b, Yangzhu Zhang^a, Yakov Kuzyakov^{c,d}, Qing Zhou^a, Tida Ge^b, Cuihong Wang^a

^a College of Resources & Environment, Hunan Agricultural University, Changsha 410128, China

^b Key Laboratory of Agro-ecological Processes in the Subtropical Region, Institute of Subtropical Agriculture, the Chinese Academy of Sciences, Changsha 410125, China

^c Department of Soil Science of Temperate Ecosystems, University of Göttingen, Göttingen 37077, Germany

^d Institute of Environmental Sciences, Kazan Federal University, Kazan 420008, Russia

ARTICLE INFO

Article history:

Received 27 November 2014

Received in revised form

14 May 2015

Accepted 16 May 2015

Available online 2 June 2015

Keywords:

Subsoil carbon

Soil organic matter

Permanganate oxidation

Chloroform fumigation

Land degradation

Deforestation

Agricultural management practices

ABSTRACT

Topsoil carbon (C) stocks are known to decrease as a consequence of the conversion of natural ecosystems to plantations or croplands; however, the effect of land use change on subsoil C remains unknown. Here, we hypothesized that the effect of land use change on labile subsoil organic C may be even stronger than for topsoil due to upward concentration of plantations and crops root systems. We evaluated soil labile organic C fractions, including particulate organic carbon (POC) and its components [coarse POC and fine POC], light fraction organic carbon (LFOC), readily oxidizable organic carbon, dissolved organic carbon (DOC) and microbial biomass down to 100 cm soil depth from four typical land use systems in subtropical China. Decrease in fine root biomass was more pronounced below 20 cm than in the overlying topsoil (70% vs. 56% for plantation and 62% vs. 37% for orchard, respectively) driving a reduction in subsoil labile organic C stocks. Land use changes from natural forest to Chinese fir plantation, Chinese chestnut orchard, or sloping tillage reduced soil organic C stocks and that of its labile fractions both in top and subsoil (20–100 cm). POC reduction was mainly driven by a decrease in fine POC in topsoil, while DOC was mainly reduced in subsoil. Fine POC, LFOC and microbial biomass can be useful early indicators of changes in topsoil organic C. In contrast, LFOC and DOC are useful indicators for subsoil. Reduced proportions of fine POC, LFOC, DOC and microbial biomass to soil organic C reflected the decline in soil organic C quality caused by land use changes. We conclude that land use changes decrease C sequestration both in topsoil and subsoil, which is initially indicated by the labile soil organic C fractions.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Land use and land use changes (LULUC) in tropical and subtropical areas, including forest conversion, capitalized agricultural intensification and animal husbandry expansion, represent major anthropogenic contributions to greenhouse gas emissions (Harris et al., 2012; IPCC, 2013). Tropical and subtropical Asia concentrates the fastest and most dramatic LULUC in the world, mainly as consequences of rapid agricultural expansion and increasing population pressure (Houghton, 2002; Carlson et al., 2012). The average rate of deforestation in tropical Asia during the 1990s reached up to

$5.6 \times 10^6 \text{ ha yr}^{-1}$, resulting in the emission of 1.0 Pg C yr^{-1} into the atmosphere (Houghton, 2002).

Tropical and subtropical aboveground biomass has received much research attention because these regions are highly productive with dense C stocks (Lewis et al., 2009; Huntingford et al., 2013). However, comprehensively studies regarding underground soil organic C (SOC) content and fractions, lability and response to land use change remain scarce.

Highly weathered tropical and subtropical soils present the deepest profiles and largest volumes among soils worldwide, accounting for nearly half of the global soil C stock in the top 3 m of soil (Richter and Markewitz, 1995; Jobbágy and Jackson, 2000). Unfortunately, most studies on the effect of LULUC on SOC have focused on the topsoil layer (0–20 cm) being the layer of soil containing the highest levels of SOC and the greatest microbial

* Corresponding author. Tel./fax: +86 731 84617803.
E-mail address: shenghao82@163.com (H. Sheng).